

# Spider Bites

## What is it?

A bite from any spider.

Most spider bites cause only minor injury but there are a few spider species that can be dangerous.

## Plan of Action:

### Seek medical care immediately if:

- You were bitten by a dangerous spider, such as a black widow or a brown recluse
- You're unsure whether the bite was from a dangerous spider
- You have severe pain, abdominal cramping or a growing wound at the bite site
- You're having problems breathing or swallowing
- The area of the sore has spreading redness or red streaks.

### To take care of a spider bite:

- Clean the wound with mild soap and water. Then apply an antibiotic ointment three times a day to help prevent infection.
- Apply a cool compress over the bite for 15 minutes each hour. Use a clean cloth dampened with water or filled with ice. This helps reduce pain and swelling.
- If possible, elevate the affected area.

- Take an over-the-counter pain reliever as needed.
- If the wound is itchy, an antihistamine, such as diphenhydramine (Benadryl) or certirizine (Zyrtec) might help.

For pain and muscle spasms, your doctor might prescribe pain medicine, muscle relaxants or both. You might also need a tetanus shot.

## Dangerous Spiders:

Black widow spider

### Black Widow spiders

You can usually identify a black widow spider by the red hourglass marking on its belly. In the United States, this spider is more common in the South. It's also found in Europe.

#### Signs and symptoms of a black widow spider bite can include:

- Redness, pain and swelling
- Severe abdominal rigidity or cramping
- Nausea, vomiting, tremors or sweating

Brown recluse spider

### Brown recluse spider

The brown recluse spider has a violin-shaped marking on its back, but this mark can be hard to see. This spider is commonly found in the southern half of the United States and in South America, where it is known as the brown spider.

#### Signs and symptoms of a brown recluse spider bite can include:

- At first, a mild pain
- Fever, chills and body aches

- A deep blue or purple area around the bite, which may develop a red ring around it

---

Revision #2

Created 6 March 2023 21:01:55

Updated 7 April 2023 15:19:15