

Anaphylaxis (Allergic Reaction)

What is it?

A life-threatening allergic reaction (anaphylaxis) can cause shock, a sudden drop in blood pressure and trouble breathing. In people who have an allergy, anaphylaxis can occur minutes after exposure to a specific allergy-causing substance (allergen). In some cases, there may be a delayed reaction, or anaphylaxis may occur without an obvious trigger.

Symptoms:

- Skin reactions, including hives, itching, and skin that becomes flushed or changes color
- Swelling of the face, eyes, lips or throat
- Narrowing of the airways, leading to wheezing and trouble breathing or swallowing
- A weak and rapid pulse
- Nausea, vomiting or diarrhea
- Dizziness, fainting or unconsciousness

Possible Triggers:

- Medications
- Latex
- Foods such as peanuts, tree nuts, fish and shellfish
- Insect stings from bees, yellow jackets, wasps, hornets and fire ants

Plan of Action:

- Immediately call 911 or your local medical emergency number.
- Ask if the person is carrying an epinephrine autoinjector (EpiPen, Auvi-Q, others) to treat an allergic attack.
- If the person needs to use an autoinjector, ask whether you should help inject the medication. This is usually done by pressing the autoinjector against the person's thigh.
- Have the person lie face up and be still.
- Loosen tight clothing and cover the person with a blanket. Don't give the person anything to drink.
- If there's vomiting or bleeding from the mouth, turn the person to the side to prevent choking.
- If there are no signs of breathing, coughing or movement, begin CPR. Do uninterrupted chest presses — about 100 every minute — until paramedics arrive.

- Get emergency treatment even if symptoms start to improve. After anaphylaxis, it's possible for symptoms to start again (recur). Monitoring in a hospital for several hours is usually necessary.

If you're with someone having symptoms of anaphylaxis, don't wait to see whether symptoms get better. Seek emergency treatment right away. In severe cases, untreated anaphylaxis can lead to death within half an hour.

An antihistamine pill, such as diphenhydramine (Benadryl), isn't enough to treat anaphylaxis. These medications can help relieve allergy symptoms, but they work too slowly in a severe reaction.

Revision #6

Created 3 March 2023 18:55:32

Updated 7 April 2023 15:18:34